

The crystallography of color superconductivity*

Jeffrey A. Bowers^{† a}, Krishna Rajagopal^a

^aCenter for Theoretical Physics, MIT, Cambridge, MA 02139, U.S.A.

We describe the crystalline phase of color superconducting quark matter. This phase may occur in quark matter at densities relevant for compact star physics, with possible implications for glitch phenomena in pulsars [1]. We use a Ginzburg-Landau approach to determine that the crystal has a face-centered-cubic (FCC) structure [2]. Moreover, our results indicate that the phase is robust, with gaps, critical temperature, and free energy comparable to those of the color-flavor-locked (CFL) phase [2]. Our calculations also predict “crystalline superfluidity” in ultracold gases of fermionic atoms [3].

Cold dense quark matter is a color superconductor [4]. At asymptotically high densities, the ground state of QCD with quarks of three flavors (u , d , and s) is the color-flavor-locked (CFL) phase [5]. This phase features a BCS condensate of Cooper pairs of quarks that includes ud , us , and ds pairs. At intermediate densities, however, the CFL phase can be disrupted by any flavor asymmetry (such as a chemical potential difference or a mass difference) that would, in the absence of pairing, separate the Fermi surfaces. In the absence of pairing, electrically neutral bulk quark matter with $m_{u,d} = 0$ and $m_s \neq 0$ features a nonzero electron density ($\mu_e \approx m_s^2/4\mu$) and three disparate quark Fermi momenta: $p_F^d \approx p_F^u + m_s^2/4\mu$, $p_F^s \approx p_F^u - m_s^2/4\mu$. (Note that decreasing μ enhances the flavor disparity.) Accounting for pairing effects modifies this picture: starting in the CFL phase at large μ , as we decrease μ the CFL phase remains “rigid” [6], with coincident quark Fermi surfaces and no electrons, until either hadronization or a first-order unlocking transition, whichever comes first. Unlocking occurs at $\mu \approx m_s^2/4\Delta_0$, where Δ_0 is the CFL gap. Its value and that of m_s are density dependent and sufficiently uncertain that we do not know whether unlocking occurs before hadronization. Here, we pursue the consequences of assuming that unlocking occurs first.

In quark matter below the unlocking transition, pairing can still occur. One option is single-flavor pairing (uu , dd , ss), but these $J = 1$ condensates have very small gaps [8]. Crystalline color superconductivity is more robust [1, 2]. We propose that unlocked quark matter is in the crystalline phase, which therefore occupies the window in the QCD phase diagram between the CFL and hadronic phases (Fig. 1). The crystalline phase has the unique virtue of allowing pairing between quarks with unequal Fermi surfaces. It was originally described by Larkin, Ovchinnikov, Fulde, and Ferrell (LOFF) [9] as a novel pairing mechanism for an electron superconductor with a Zeeman splitting between

[†]Speaker.

*We thank the Kavli Institute for Theoretical Physics in Santa Barbara for support and hospitality. Research supported in part by DOE cooperative research agreement #DF-FC02-94ER40818, by NSF Grant PHY99-07949, and (that of JAB) by a KITP Graduate Fellowship and DOD NDSEG Fellowship.




Figure 1. Schematic QCD phase diagram with quark-gluon plasma, crystalline (XTAL) and CFL color superconducting phases, the latter separated by the unlocking transition.




Figure 2. A unit cell of the FCC crystal, with contours where the gap function is positive (black), zero (gray), and negative (white).

spin-up and spin-down Fermi surfaces, neglecting all orbital effects of the magnetic perturbation. Quark matter is a more natural setting for the LOFF phase, as it features a “flavor Zeeman effect” with no orbital complications. Cooper pairs in the LOFF phase have nonzero total momentum: a quark with momentum $\mathbf{p}$ is paired with a quark with momentum $-\mathbf{p} + 2\mathbf{q}$. The magnitude $|\mathbf{q}| \equiv q_0$ is determined by the separation between Fermi surfaces while the direction(s) $\hat{\mathbf{q}}$ is (are) chosen spontaneously. Each quark in a pair resides near its respective Fermi surface, so pairs can form at low cost and a condensate results. For a given $\mathbf{q}$, the quarks that pair are only those in “pairing rings”, one on each Fermi surface; these circular bands are antipodal to each other and perpendicular to $\mathbf{q}$.

If each Cooper pair in the condensate carries the same total momentum $2\mathbf{q}$, then in position space the condensate varies like a plane wave: $\langle \psi(\mathbf{x})\psi(\mathbf{x}) \rangle \sim \Delta \exp(2i\mathbf{q} \cdot \mathbf{x})$ meaning that translational and rotational symmetry are broken. If the system is unstable to the formation of a single plane-wave condensate, then we expect that a superposition of multiple plane waves is more favorable: $\langle \psi(\mathbf{x})\psi(\mathbf{x}) \rangle \sim \sum_{\mathbf{q}, |\mathbf{q}|=q_0} \Delta_{\mathbf{q}} e^{2i\mathbf{q} \cdot \mathbf{x}}$. Each $\Delta_{\mathbf{q}}$ corresponds to condensation of Cooper pairs with momentum $2\mathbf{q}$, i.e. another pairing ring on each Fermi surface. As we add more plane waves, we utilize more of the Fermi surface for pairing, with a corresponding gain in condensation energy. On the other hand, the rings can “interact” with each other: condensation in one mode can enhance or deter condensation in another mode. We need to compare various different choices for the set of $\hat{\mathbf{q}}$ ’s whose $\Delta_{\mathbf{q}}$ ’s are nonzero; that is, we need to compare different crystal structures. To this end, we calculate the Ginzburg-Landau free energy $\Omega(\{\Delta_{\mathbf{q}}\})$ for the LOFF phase.

We consider a simplified model with massless u and d quarks only, and introduce chemical potentials $\mu_u = \bar{\mu} - \delta\mu$, $\mu_d = \bar{\mu} + \delta\mu$. In this toy model, we vary $\delta\mu$ by hand to emulate the effect of splitting the Fermi surfaces as described above. (The LOFF phenomenon can equally well be induced by mass differences between quarks [10].) We work at zero temperature, as appropriate for compact star physics. We model the interaction between

quarks with a pointlike four-fermion interaction that mimics single gluon exchange.

At $\delta\mu = 0$, the system forms a BCS superconductor with a gap Δ_0 . As we increase $\delta\mu$, the system exhibits a “rigidity” analogous to that of the CFL phase: despite the imposed stress $\delta\mu$, the gap stays constant and the Fermi surfaces remain coincident. When $\delta\mu$ reaches a critical value $\delta\mu_1 = \Delta_0/\sqrt{2} \approx 0.707\Delta_0$, the BCS phase “breaks” and the Fermi surfaces separate (this is analogous to CFL “unlocking”). For $\delta\mu > \delta\mu_1$ a crystalline phase is possible. If we consider the simplest LOFF ansatz, a single plane wave condensate [1], we find that the LOFF state is favored in an interval $\delta\mu_1 \leq \delta\mu \leq \delta\mu_2$; at $\delta\mu_1$ there is a first-order transition from BCS to LOFF, while at $\delta\mu_2 \approx 0.754\Delta_0$ there is a second-order transition from LOFF to the normal state (unpaired quarks). Each Cooper pair has a total momentum $2\mathbf{q}$, where $|\mathbf{q}| = q_0 \approx 1.20\delta\mu$; this implies that the pairing ring on each Fermi surface has an opening angle $\psi_0 \approx 2\cos^{-1}(\delta\mu/q_0) \approx 67.1^\circ$.

Motivated by the second-order transition at $\delta\mu_2$, we write a Ginzburg-Landau free energy valid in the small order parameter limit [2]. The most general expression consistent with translational and rotational symmetry is

$$\Omega \propto \sum_{\mathbf{q}, |\mathbf{q}|=q_0} \alpha \Delta_{\mathbf{q}}^* \Delta_{\mathbf{q}} + \frac{1}{2} \sum_{\square} J(\square) \Delta_{\mathbf{q}_1}^* \Delta_{\mathbf{q}_2} \Delta_{\mathbf{q}_3}^* \Delta_{\mathbf{q}_4} + \frac{1}{3} \sum_{\diamond} K(\diamond) \Delta_{\mathbf{q}_1}^* \Delta_{\mathbf{q}_2} \Delta_{\mathbf{q}_3}^* \Delta_{\mathbf{q}_4} \Delta_{\mathbf{q}_5}^* \Delta_{\mathbf{q}_6} + \dots \quad (1)$$

The symbol $\square$ represents a set of four equal-length vectors $(\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4)$, $|\mathbf{q}_i| = q_0$, with $\mathbf{q}_1 - \mathbf{q}_2 + \mathbf{q}_3 - \mathbf{q}_4 = 0$, forming a closed figure. Similarly, the symbol $\diamond$ represents a set of six equal-length vectors $(\mathbf{q}_1, \dots, \mathbf{q}_6)$ with $\mathbf{q}_1 - \mathbf{q}_2 + \mathbf{q}_3 - \mathbf{q}_4 + \mathbf{q}_5 + \mathbf{q}_6 = 0$. The quadratic coefficient α changes sign at $\delta\mu_2$ showing the onset of the LOFF plane-wave instability: $\alpha \approx (\delta\mu - \delta\mu_2)/\delta\mu_2$. Analysis of the quadratic term shows that all modes on the sphere $|\mathbf{q}| = q_0$ become unstable at $\delta\mu < \delta\mu_2$, and that at larger $\delta\mu$ it is these modes that are the least stable. The quartic and sextic coefficients J and K characterize the interactions between modes, and thus determine the crystal structure. For candidate crystal structures with all $\Delta_{\mathbf{q}}$ ’s equal in magnitude, we can evaluate aggregate Ginzburg-Landau quartic and sextic coefficients β and γ as sums over all rhombic and hexagonal combinations of the $\mathbf{q}$ ’s: $\beta = \sum_{\square} J(\square)$, $\gamma = \sum_{\diamond} K(\diamond)$. Then for a crystal with P plane waves we obtain $\Omega(\Delta) \propto P\alpha\Delta^2 + \frac{1}{2}\beta\Delta^4 + \frac{1}{3}\gamma\Delta^6 + \dots$ and we compare crystals by calculating β and γ to find the structure with the lowest Ω .

We have analyzed many candidate crystal structures in [2]; here, we summarize the lessons learned. Recall that each $\mathbf{q}$ corresponds to a pairing ring on each Fermi surface with opening angle $\psi_0 \approx 67.1^\circ$. The first lesson is that adding more pairing rings lowers the free energy only as long as rings do not intersect. No more than nine rings can be arranged on the Fermi surface without any overlaps. The second lesson is that “regular” structures (those with nonintersecting rings whose wave vectors can be assembled into closed figures in many ways) are combinatorically favored. There are no particularly regular nine-wave structures, but there is a very regular eight-wave structure: eight wave vectors $\mathbf{q}$ pointing towards the eight corners of a cube, forming the eight shortest vectors in the reciprocal lattice of a face-centered-cubic (FCC) crystal. This FCC crystal has by far the lowest Ginzburg-Landau free energy of all the structures we considered. The gap function is $\langle \psi(\mathbf{x})\psi(\mathbf{x}) \rangle \sim \Delta [\cos \frac{2\pi}{a}(x+y+z) + \cos \frac{2\pi}{a}(x-y+z) + \cos \frac{2\pi}{a}(x+y-z) + \cos \frac{2\pi}{a}(-x+y-z)]$ with lattice constant $a = \sqrt{3}\pi/|\mathbf{q}| \approx 6.012/\Delta_0$. A unit cell is shown in Fig. 2. The effective field theory for the phonons of this crystal has recently been constructed [11].

The Ginzburg-Landau free energy for the FCC structure has coefficients β and γ that are large and negative. This guarantees that the phase transition to the crystalline phase cannot be a second-order transition at $\delta\mu = \delta\mu_2$ and must instead be a strongly first-order transition at some $\delta\mu = \delta\mu_* \gg \delta\mu_2$, at which the magnitude of the gap jumps from zero to a large nonzero value. Were we to push beyond sextic order to obtain a bounded Ginzburg-Landau free energy, the resulting calculation of the gap and ground state free energy would be quantitatively uncontrolled because the order parameter is nowhere nonzero and small. Qualitatively, the large negative coefficients indicate a robust crystalline phase, with gaps and free energy comparable to those of the BCS phase. Furthermore, because the transition to the crystalline phase occurs at $\delta\mu_* \gg \delta\mu_2$, the crystalline phase is favored over a wide $\delta\mu$ interval. Our results thus indicate that wherever unlocked quark matter occurs, it will be in the crystalline phase with an FCC crystal structure.

The power of the Ginzburg-Landau formulation is that it organizes the calculation in such a way that simple qualitative lessons about what features make a crystal structure favorable emerge. We immediately learn that, among all the possibilities, the FCC structure of Fig. 2 is the best choice. This is the first time (in either the condensed matter or the QCD incarnations of this subject) that sufficiently general three dimensional crystal structures have been compared. The structure that we find to be most favorable, although simple with hindsight, was never considered in the classic condensed matter physics papers. Unfortunately, but not unusually, the Ginzburg-Landau calculation predicts a first-order phase transition and in so doing predicts its own quantitative demise. With a qualitative understanding of what crystal structure wins (that is, with confidence in what ansatz to make) we are currently doing the variational calculation needed to determine the gap, free energy, and $\delta\mu_*$ quantitatively within our toy model. Going beyond our toy model, a full three-flavor analysis is desirable, with ud , us and ds crystalline condensates and a strange quark mass. In an atomic physics context, the FCC condensate may be detectable as a “crystalline superfluid” in an ultracold gas of fermionic atoms [3]. In an astrophysics context, with the crystal structure known a calculation of the pinning force can now proceed, in order to determine whether (some) observed pulsar glitches originate from pinning of rotational vortices at intersections of crystal nodal planes.

REFERENCES

1. M. Alford, J. A. Bowers and K. Rajagopal, Phys. Rev. **D63**, 074016 (2001).
2. J. A. Bowers and K. Rajagopal, hep-ph/0204079.
3. R. Combescot, Europhys. Lett. **55**, 150 (2001).
4. Reviews: M. Alford, hep-ph/0102047; K. Rajagopal, F. Wilczek, hep-ph/0011333; T. Schäfer, E. Shuryak, nucl-th/0010049; D. Rischke, R. Pisarski, hep-ph/0004016.
5. M. Alford, K. Rajagopal and F. Wilczek, Nucl. Phys. B **537**, 443 (1999).
6. K. Rajagopal and F. Wilczek, Phys. Rev. Lett. **86**, 3492 (2001).
7. M. Alford and K. Rajagopal, JHEP **0206**, 031 (2002).
8. T. Schäfer, Phys. Rev. **D62**, 094007 (2000).
9. A. I. Larkin and Yu. N. Ovchinnikov, Sov. Phys. JETP **20**, 762 (1965); P. Fulde and R. A. Ferrell, Phys. Rev. **135**, A550 (1964).
10. J. Kundu and K. Rajagopal, Phys. Rev. D **65**, 094022 (2002).
11. R. Casalbuoni, E. Fabiano, R. Gatto, M. Mannarelli and G. Nardulli, hep-ph/0208121.